Alcohol addiction/Quiz

The questions in the quiz refer to the article on the expert article on alcohol addiction.

women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	
Points for an incorrect answer: Ignore the questions' coefficients: Shuffle questions 1 Which answer regarding the prevalence of alcohol dependence is correct? In Germany, men drink alcohol more frequently and in larger quantities than women. Alcohol dependence is clearly more common among men, alcohol abuse among women. Alcohol abuse and dependence occur most frequently in old age. The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	
Points for an incorrect answer: Ignore the questions' coefficients: Shuffle questions 1 Which answer regarding the prevalence of alcohol dependence is correct? In Germany, men drink alcohol more frequently and in larger quantities than women. Alcohol dependence is clearly more common among men, alcohol abuse among women. Alcohol abuse and dependence occur most frequently in old age. The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	
Points for an incorrect answer: Ignore the questions' coefficients: Shuffle questions 1 Which answer regarding the prevalence of alcohol dependence is correct? In Germany, men drink alcohol more frequently and in larger quantities than women. Alcohol dependence is clearly more common among men, alcohol abuse among women. Alcohol abuse and dependence occur most frequently in old age. The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	
Points for an incorrect answer: Ignore the questions' coefficients: Shuffle questions 1 Which answer regarding the prevalence of alcohol dependence is correct? In Germany, men drink alcohol more frequently and in larger quantities than women. Alcohol dependence is clearly more common among men, alcohol abuse among women. Alcohol abuse and dependence occur most frequently in old age. The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	
Ignore the questions' coefficients: Shuffle questions 1 Which answer regarding the prevalence of alcohol dependence is correct? In Germany, men drink alcohol more frequently and in larger quantities than women. Alcohol dependence is clearly more common among men, alcohol abuse among women. Alcohol abuse and dependence occur most frequently in old age. The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	t added for a correct answer: 1
Shuffle questions 1 Which answer regarding the prevalence of alcohol dependence is correct? In Germany, men drink alcohol more frequently and in larger quantities than women. Alcohol dependence is clearly more common among men, alcohol abuse among women. Alcohol abuse and dependence occur most frequently in old age. The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	ts for an incorrect answer:
Which answer regarding the prevalence of alcohol dependence is correct? In Germany, men drink alcohol more frequently and in larger quantities than women. Alcohol dependence is clearly more common among men, alcohol abuse among women. Alcohol abuse and dependence occur most frequently in old age. The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	re the questions' coefficients:
Which answer regarding the prevalence of alcohol dependence is correct? In Germany, men drink alcohol more frequently and in larger quantities than women. Alcohol dependence is clearly more common among men, alcohol abuse among women. Alcohol abuse and dependence occur most frequently in old age. The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	uffle questions
 In Germany, men drink alcohol more frequently and in larger quantities than women. Alcohol dependence is clearly more common among men, alcohol abuse among women. Alcohol abuse and dependence occur most frequently in old age. The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women. 	
 Alcohol dependence is clearly more common among men, alcohol abuse among women. Alcohol abuse and dependence occur most frequently in old age. The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women. 	th answer regarding the prevalence of alcohol dependence is correct?
 Alcohol abuse and dependence occur most frequently in old age. The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women. 	Germany, men drink alcohol more frequently and in larger quantities than women.
 The male dominance of dependence already develops in adolescence. Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women. 	cohol dependence is clearly more common among men, alcohol abuse among women.
Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	cohol abuse and dependence occur most frequently in old age.
 Which of the following statements about the pathophysiology is not true? Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women. 	ne male dominance of dependence already develops in adolescence.
 Women show an increased vulnerability to the toxic effects of alcohol. At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women. 	
 At similar alcohol levels, men react more aggressively on average than women. Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women. 	
Alcohol consumption increases the risk of cancer of the mouth, pharynx and oesophagus, women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	
women. Alcohol consumption is associated, among other things, with spontaneous miscarriages or development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	
 development of a disorder from the range of foetal alcohol syndrome. Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women. 	·
 Which statement regarding possible therapies is not true? Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women. 	cohol consumption is associated, among other things, with spontaneous miscarriages or the evelopment of a disorder from the range of foetal alcohol syndrome.
 Men seem to benefit more from clear structures (such as in Alcoholics Anonymous). Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women. 	
Women are more likely to need groups where emotions are worked through and skills succonfidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	
confidence or self-efficacy are practised for treatment success. For women, gender-homogeneous groups are more recommended, as topics relevant to the discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	· · · · · · · · · · · · · · · · · · ·
discussed better this way. For men, social support is a factor that has a stronger influence on both the uptake of the success of treatment than it does for women.	omen are more likely to need groups where emotions are worked through and skills such as self- infidence or self-efficacy are practised for treatment success.
success of treatment than it does for women.	or women, gender-homogeneous groups are more recommended, as topics relevant to them can be scussed better this way.
C.h.v.'s	or men, social support is a factor that has a stronger influence on both the uptake of therapy and the occess of treatment than it does for women.
Submit	mit